

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/326985141>

International Journal of Fisheries and Aquatic Studies 2016; 4(5): 233–236

Length–Weight relationships of some economic freshwater fishes of Nwaniba River, Southeast Nigeria

Article · August 2018

CITATIONS

0

READS

71

3 authors, including:



Imeh Esenowo

University of Uyo

24 PUBLICATIONS 77 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



soil arthropods sampling using different sampling techniques [View project](#)



Toxicological Effects on different concentrations of commonly used bleach on the Survival of African Clariid Mud Catfish (*Clarias gariepinus*) Fingerlings [View project](#)



International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.549

IJFAS 2016; 4(5): 233-236

© 2016 IJFAS

www.fisheriesjournal.com

Received: 01-07-2016

Accepted: 01-08-2016

Esenowo IK

Department of Zoology,
University of Uyo, Akwa Ibom
State, Nigeria

Ogidiaka E

Department of Science and
Technology, Fisheries and
Fisheries Technology Option,
Delta State School of Marine
Technology, Burutu, Delta
State, Nigeria

Brownson II

Department of Fisheries and
Aquaculture, University of Uyo,
Akwa Ibom State, Nigeria

Length-Weight relationships of some economic freshwater fishes of Nwaniba River, Southeast Nigeria

Esenowo IK, Ogidiaka E and Brownson II

Abstract

The length-weight relationships of some economic freshwater fishes were analyzed in Nwaniba River, Southern Nigeria. Fish specimens were procured from middlemen at the landing site from January to June, 2014. The mean total length and weight were 15.76 cm and 78.39g respectively. The highest (3.512) and lowest (1.503) “a” value was recorded for *Hemichromis fasciatus* and *Oreochromis niloticus* respectively, while the “b” value ranged from 2.78 and 3.54. The mean condition factor ranged from 1.818 and 2.351. The result obtained in this study showed both negative and positive allometric growth pattern.

Keywords: Length-weight, economic freshwater fishes, condition factor, Nwaniba River

1. Introduction

Fish, especially those in tropical and sub-tropical water system are known to experience growth fluctuations which are due to factors such as changes in environmental parameters and food composition [27, 1, 25]. Apart from changes in these environmental characteristics, the morphometric characteristics are also used to assess growth fluctuation on fish populations. In this regard, it is common to use measurements such as body-length, body depth, head length, eye diameter, jaw length of fishes etc, to not only assess the fish habitat peculiarities and ecological criteria in water bodies, but to also measure discreteness and relationships among various taxonomic categories [19, 26].

Length and weight data of fish are useful parameters in estimating the length and age structures, population dynamic, growth, mortality rates and well-being of the fish [16, 17, 4, 22, 18]. They are also important tools used to obtain information on length frequency distribution [3, 12], fish condition for stock assessment [28, 1] and management of the fish population [31, 7, 13].

Condition factor in fisheries is crucial as a quantitative parameter because the heavier the fish species of a given length, the better the physiological condition of the aquatic organism [6, 33]. It is also an index to understand the life-cycle of a fish by referring to the coefficient values derived from its length-weight relationship data [30]. However, the condition factor of fish is strongly affected by biotic and abiotic environmental variables [20, 29, 4]. Recent studies on the fish fauna in Nwaniba River have been conducted on species richness and diversity of ichthyofaunal and ornamental fish species by some researchers [24, 8, 32, 9], but little or no work on the Length-weight Relationship of the fish species in the study area of Nwaniba River, hence, the need for the present study to provide information on an estimated average weight of fish species of a given length, as well as using this data to assess the relative wellbeing of some fish population in this river.

2. Materials and Methods

2.1 The Study area

Nwaniba River is located between latitude 5°2'51" N and longitude 5°2'41" E of Southeast Nigeria. It is one of the freshwater rivers in Uruan Local Government Area of Akwa Ibom State, Nigeria which flows from Itam River in Itu Local Government Area through Mbiakong River. The depth of the river ranged from 0.2 - 8.4m with a mean value of 3.75 ±0.57m. The annual rainfall is about 2500mm with temperature of 32°C and relative humidity of 75%. The study area comprises dry (November-March) and rainy (April-October) seasons.

Correspondence

Esenowo IK

Department of Zoology,
University of Uyo, Akwa Ibom
State, Nigeria

The bank of the river is mostly covered with grasses while other portions have swamp vegetation. The occupation of the local people is fishing, farming, trading, saw-milling and canoe construction. Artisanal fishermen within the river

mainly exploit the fisheries using wooden dug-out canoes ranging from 5m to 50m long which serves as a means of transportation.




Fig 1: Map showing Nwaniba River

2.2 Collection of samples

Fish specimens were procured from middlemen at the landing site from January to June, 2014. The fishermen used a wide range of fishing gear such as hook and line, long line, cast nets, gill nets and traps. Fish specimens were randomly chosen and identified using keys and descriptions by [15]. The length-weight relationship was calculated using the least square regression on log transformation of the equation:

$$\log W = \log a + b \log L$$

Where W = Weight, TL = Total Length, a = exponent describing the rate of change of weight with length

(intercept), b = Slope

The Condition factor, k was calculated using this formula; $K = 100W/L^3$ while the Pearson correlation coefficient was used to determine the strength of relationship between the weight and length of each fish.

3. Results

The results of the measured standard lengths (TL) and body weights (TW) of the fishes examined are presented in Table 1. The values of the regression coefficients a & b and the condition factors obtained are presented in Table 2.

Table 1: The Standard lengths and body weight of fishes examined

Fish	Total Length (TL) (cm)		Total Weight (TW) (g)	
	Size Range	Mean	TW Range	Mean
<i>Chromidotilapia guntheri guntheri</i>	14.4-18.2	16.18±1.15	59.9-105.37	80.4±15.86
<i>Coptodon guineensis</i>	8.3-20.0	14.71±2.23	10.05-157.89	74.69±29.90
<i>Coptodon zilli</i>	13.5-21.3	16.03±3.56	41.62-179.52	83.83±64.21
<i>Hemichromis fasciatus</i>	13.3-16.9	14.60±1.16	42.2-112.97	63.04±26.20
<i>Oreochromis niloticus</i>	8.0-19.7	15.40±3.04	9.98-132.85	67.89±31.55
<i>Sarotherodon melanotheron</i>	16.6-19.5	17.68±1.33	79.2-134.50	100.5±24.79

Table 2: The regression coefficient and condition factors of fishes examined

Fish	a	b	r	K
<i>Chromidotilapia guntheri guntheri</i>	1.514	2.82	0.9753	1.891
<i>Coptodon guineensis</i>	1.642	2.99	0.9661	2.351
<i>Coptodon zilli</i>	1.915	3.14	0.9989	2.047
<i>Hemichromis fasciatus</i>	3.512	3.54	0.9742	1.985
<i>Oreochromis niloticus</i>	1.503	2.78	0.9913	1.856
<i>Sarotherodon melanotheron</i>	1.991	3.20	0.9919	1.818

The exponent (b) values for *Chromidotilapia guntheri guntheri*, *Coptodon guineensis* and *Oreochromis niloticus*

were less than 3, indicating negative allometric growth pattern while the exponent (b) values for *Coptodon zilli*, *Hemichromis fasciatus* and *Sarotherodon melanotheron* were greater than 3 indicating positive allometric growth. The correlation coefficients values (r) ranged between 0.96617 and 0.99895 in all six fishes and this showed a high degree of positive correlation between the SL and BWT.

4. Discussions

The sizes of *S. melanotheron* and *C. guntheri guntheri* examined were larger than those of *C. guineensis*, *C. zilli*, *H. fasciatus* and *O. Niloticus* in this study. The variation in the length and weight could probably be attributed to their faster

growth rates and voracious predatory and carnivorous feeding habits [14, 2]. Oni *et al.*, [23] also reported that feeding and growth were the major factors responsible for the size variations of these fish species in River Galma, Zaria Dam. However, it was observed that the large size fishes were adult, probably with full-laden stomachs or with matured eggs which are due to gear non-selective techniques resulting in samples ranging from immature to fully matured fishes [11].

Chromidotilapia guntheri had a mean condition factor of 1.891 despite having a mean body length and mean body weight of 16.18±1.15 cm and 80.4±15.86 respectively. Anene [4] reported a relatively lower condition factors for relatively large sizes, while relatively higher condition factors were recorded for rather smaller fish. The reason for this result may likely be that adults spend a part of their energy in reproduction, hence the higher condition factor for larger fishes as reported by [5].

The *b* value for *S. melanotheron* in this result of $b = 3.20$ varies with juveniles of *S. melanotheron* and adults *S. melanotheron* of $b = 2.85$ and $b = 2.87$ respectively from Lake Nokoué and Ahémé as reported by [21]. Also, the “*b*” values obtained during this study for the same species is higher than 2.8 as reported by [5] in Eleiyele Lake and same with 2.2 by [24] in Ntak Inyang stream, Ikpa River. However, the values of the regression coefficient (*b*) for all the fish species for the entire period of study were significantly different ($p > 0.05$) from 3.0 which is an indication that they exhibited different growth pattern, ranging from positive allometric to negative allometric. The growth pattern observed in this work implies that the fish species may be longer than its weight or weight increases faster than its length. This may be attributed to over-fishing by the natives owing to easy accessibility to the sampling stations, hence making it difficult for the species to grow to a sizeable population. Similar report by [10] show *Sarotherodon galilaeus* exhibiting allometric growth pattern in Opa Reservoir.

The mean condition factors of 1.81 recorded is lower than 4.66 and 4.45 for *S. melanotheron* as reported by [21] in Lakes Nokoué and Ahémé respectively, but similar with *k* value as reported by [5] in Eleiyele Lake, Southwestern Nigeria. According to [6], *k* value for fish must range from 2 - 4 and this is similar with the *k* values ranging from 2.78- 3.54 in this study. This suggests that the condition of some fish from Nwaniba River only favour *Coptodon guineensis* and *Coptodon zilli* species most especially.

5. Conclusion

The predominant growth pattern exhibited by the sampled fish species was both negative and positive allometry growth pattern and this could be due to over-fishing by the native’s community. The mean condition factor for all species investigated was greater than 1, which indicates that the fish were above average condition in the river. This study therefore, provides baseline information on the length-weight relationship of some economic fishes in Nwaniba River that will be useful in fisheries resource management of the water body.

6. References

1. Abowei AFN, Davies OA, Eli AA. Study of the length-weight relationship and condition factor of five fish species from Nkoro River, Niger Delta, Nigeria, Current Research Journal of Biological Sciences. 2009; 1(3):94-95.
2. Abowei JFN, Hart AI. Some morphometric parameters of ten finfish species from the lower Nun River, Niger Delta, Nigeria. Res. J. Biol. Sci. 2009; 4(3):282-288.
3. Anderson RO, Gutreuter SJ. Length, weight, and associated structural indices, In Nielsen, L., Johnson, D. (Eds.) Fisheries Techniques American Fisheries Society, Bethesda, Maryland, 1983, 284-300.
4. Anene A. Condition factor of four cichlid species of a man-made lake in Imo State, Southeastern Nigeria, Turkish Journal of Fisheries and Aquatic Sciences. 2005; 5:43-47.
5. Ayoade AA, Ikulala AOO. Length weight relationship, condition factor and stomach contents of *Hemichromis bimaculatus*, *Sarotherodon melanotheron* and *Chromidotilapia guentheri* (Perciformes: Cichlidae) in Eleiyele Lake, Southwestern Nigeria. Revista de Biologia Tropical. 2007; 55(3-4):969-977.
6. Bagenal TB, Tesch AT. Conditions and Growth Patterns in Fresh Water Habitats. Blackwell Scientific Publications, Oxford, 1978, 75-89.
7. Blackwell BG, Brown ML, Willis DW. Relative weight (*W_r*) status and current use in fisheries assessment and management, Reviews in Fisheries Science. 2000; 8(1):1-44.
8. Ekpo IE, Udo MT, Usip PL. Seasonality and size variation of fish species in Nwaniba, Ikpa River Southeast Nigeria. Elixir Agriculture. 2012; 51:11043-11050
9. Ekpo IE. Ornamental fish species potentials of Ikpa River in Akwa Ibom State, Nigeria. Journal of Biology, Agriculture and Healthcare. 2013; 3(6):61-66.
10. Fawole OO, Arawomo GA. Fecundity of *Sarotherodon galilaeus* in the Opa reservoir Ile Ife, Nig. J. Sci. Res. 1999; 4(1):107-111.
11. Frota LO, Costa PAS, Braga AC. Length-weight relationships of marine fishes from the central Brazilian coast. NAGA, World Fish Center Quarterly. 2004; 27:20-26.
12. Gayanilo FC, Pauly D. FAO ICLARM stock assessment tools (FISAT)”, References Manual, FAO Computerized Information Series (Fisheries). 1997; 8:262.
13. Haimovici M, Velasco G. Length-weight relationships of marine fishes from southern Brazil. NAGA, The ICLARM Quarterly. 2000; 23(1):19-23.
14. Idodo-Umeh G. The feeding ecology of Mochokid species in River Ase, Niger Delta, Nigeria. Tropical Freshwater Biology. 2005; 14:71-93.
15. Idodo-Umeh G. Freshwater Fishes of Nigeria. (Taxonomy, Ecological Notes, Diet and Utilization). Idodo-Umeh Pub. Ltd, Benin-City, Nigeria, 2003, 232.
16. Kohler N, Casey J, Turner P. Length-weight relationships for 13 species of sharks from the western North Atlantic, Fisheries Bulletin. 1995; 93:412-418.
17. Krause J, Jean-Guy J, Brown D. Body length variation within multi-species fish shoals: the effects of shoal size and number of species, *Oecologia*. 1998; 114:67-72.
18. Koffi BK, Berté S, Koné T. Length-weight Relationships of 30 Fish Species in Aby Lagoon, Southeastern Côte d’Ivoire. Current Research Journal of Biological Sciences. 2014; 6(4):173-178, 2014.
19. Lalèyè PA. Length-weight and length-length relationships of fishes from the Ouémé River in Bénin (West Africa). Journal of Applied Ichthyology. 2006; 22:330-333.

20. Luff RM, Bailey GN. Analysis of size changes and incremental growth structures in African catfish *Synodontis schall* (schall) from Tell el-Amarna, Middle Egypt, Journal of Archaeological Science. 2000; 27:821-835.
21. Niyonkuru C, Laleye P. A Comparative Ecological Approach of the Length–Weight Relationships and Condition Factor of *Sarotherodon Melanotheron* Rüppell, 1852 and *Tilapia Guineensis* (Bleeker 1862) in Lakes Nokoué and Ahémé (Bénin, West Africa) International Journal of Business, Humanities and Technology, 2012, 2(3).
22. Ndimele PE, Kumolu-Johnson CA, Aladetohun NF, Ayorinde OA. Length-weight relationship, condition factor and dietary composition of *Sarotherodon melanotheron*, Rüppell, 1852 (Pisces: cichlidae) in Ologe Lagoon, Lagos, Nigeria. Agriculture and Biology Journal of North America. 2010; 1(4):584-590.
23. Oni SK, Olayemi JY, Adegboye JD. The comparative physiology of three ecologically (Rupel). *Synodontis schall*. Block and Schneider and *Tilapia zilli* (Gervais). Journal of Fish Biology. 1983; 22:105-109.
24. Onuoha GC, Ekpo IE, Chude LA, Isangedighi IA. Composite preliminary ichthyofaunal survey of Ntak Nyang stream, Ikpa River, Nigeria. Nigerian Journal of Agriculture, Food & Environment. 2010; 6(1&2):82-89.
25. Obasohan EE, Obasohan EE, Imasuen JA, Isidahome CE. Preliminary studies of the length-weight relationships and condition factor of five fish species from Ibiekuma stream, Ekpoma, Edo state, Nigeria. Journal of Agricultural research and development. 2012; 2(3):61-69.
26. Omoniyi I, Agbon A, Sodunke SA. Effects of Lethal and Sub-lethal concentrations of Tobacco (*Nicotiana tobaccum*), leaf dust extraction on weight and haematological changes in *Clarias gariepinus* (Burchell). J. Applied Sci. Environ. Man. 2002; 6:37-41.
27. Panfili J, Mbow A, Durand JD, Diop K, Diouf K, Ndiaye P *et al.* Influence of salinity on life-history traits of the West-African black-chinned tilapia (*Sarotherodon melanotheron*): comparison between the Gambia and Saloum estuaries. Aquatic Living Resources. 2004; 17:65-74.
28. Petrakis G, Stergiou KI. Weight-length relationship for 33 fish species in Greek waters. Fish. Res. 1995; 21:465-469.
29. Saliu JK. “Observation on the Condition Factor of *Brycinus nurse* (Pisces: Cypriniformes, Characidae) from Asa Reservoir, Ilorin, Nigeria”, Tropical Freshwater Biology. 2001; 10:9-17.
30. Schneider JC, Laarman PW, Gowing H. Length-weight relationships, In Schneider, James C. (Ed.). Manual of fisheries survey methods II: with periodic updates. Michigan Department of Natural Resources, Fisheries Special Report 25, 2000: Ann Arbor.
31. Sparre P, Venema SC. Introduction to Tropical Fish Stock Assessment, Part 1: Manual. FAO Fisheries Technical Paper. 1998; 306(1):433.
32. Udo IU. Taxonomic composition, diversity and abundance of the ichthyofaunal assemblage of Iba-Oku stream, Ikpa River, Nigeria. International Journal of Zoological Research. 2012; 8:71-80
33. Weatherley AH, Gill HS. The biology of fish growth, London, 1987, 443.